

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

AD-A208 332

DULE

1b. RESTRICTIVE MARKINGS

3 DISTRIBUTION/AVAILABILITY OF REPORT

Approved for public release;
distribution is unlimited

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

NMRI 89-17

5. MONITORING ORGANIZATION REPORT NUMBER(S)

6a. NAME OF PERFORMING ORGANIZATION
Naval Medical Research6b. OFFICE SYMBOL
(If applicable)7a. NAME OF MONITORING ORGANIZATION
Naval Medical Command

6c. ADDRESS (City, State, and ZIP Code)

Bethesda, Maryland 20814-5055

7b. ADDRESS (City, State, and ZIP Code)

Department of the Navy
Washington, D.C. 20372-51208a. NAME OF FUNDING/SPONSORING
ORGANIZATION Naval Medical8b. OFFICE SYMBOL
(If applicable)

Research and Development Command

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

8c. ADDRESS (City, State, and ZIP Code)

Bethesda, Maryland 20814-5055

10. SOURCE OF FUNDING NUMBERS

PROGRAM
ELEMENT NO.
61102APROJECT
NO.
3M161102BS13TASK
NO.
AK-111WORK UNIT
ACCESSION NO.
DA313955

11. TITLE (Include Security Classification)

Human lymphocyte proliferative response to a sporozoite T cell epitope correlates with
resistance to falciparum malaria

12. PERSONAL AUTHOR(S)

Hoffman SL, Oster CN, Mason C, Beier JC, Sherwood J,
Chulay JD

Ilou WR, Mugambi M,

13a. TYPE OF REPORT

journal article

13b. TIME COVERED

FROM TO

14. DATE OF REPORT (Year, Month, Day)

1989

15. PAGE COUNT

4

16. SUPPLEMENTARY NOTATION

Reprinted from: The Journal of Immunology, Feb 15, 1989 Vol.142(4) pp. 1299-1303

17. COSATI CODES

FIELD	GROUP	SUB-GROUP

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

Animal; Antigenic Determinants; Antigens, Protozoan;
Lymphocyte Transformation; Malaria; Plasmodium Falciparum;
T Lymphocytes

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

DTIC
ELECTE
MAY 25 1989
S D

20 DISTRIBUTION/AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION

Unclassified

22a. NAME OF RESPONSIBLE INDIVIDUAL

Phyllis Blum, Information Services Division

22b. TELEPHONE (Include Area Code)

202-295-2188

22c. OFFICE SYMBOL

ISD/ADMIN/NMRI

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted.

All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE
UNCLASSIFIED

HUMAN LYMPHOCYTE PROLIFERATIVE RESPONSE TO A SPOROZOITE T CELL EPITOPE CORRELATES WITH RESISTANCE TO FALCIPARUM MALARIA¹

STEPHEN L. HOFFMAN,^{2*} CHARLES N. OSTER,¹ CARL MASON,¹ JOHN C. BEIER,³
JAMES A. SHERWOOD,¹ W. RIPLEY BALLOU,¹ MUTUMA MUGAMBI,⁴ AND
JEFFREY D. CHULAY¹

From the ¹Infectious Diseases Department, Naval Medical Research Institute, Bethesda, MD 20814-5055; ²Department of Immunology, Walter Reed Army Institute of Research, Washington, D.C. 20307-5100; ³Kenya Medical Research Institute, United States Army Medical Research Unit Kenya, Nairobi, Kenya, and ⁴Kenya Medical Research Institute, Nairobi, Kenya

To identify vaccine relevant T cell epitopes on the circumsporozoite (CS) protein of *Plasmodium falciparum*, the lymphocyte proliferative responses to 10 CS protein derived peptides were studied in 28 adult Kenyans, and correlated with resistance to malaria. Eight peptides, six of which were not overlapping, induced proliferation of lymphocytes from one to five volunteers, suggesting either genetic restriction of response to each of the T epitopes, or dominance of some T sites on the immunizing sporozoites. The 28 volunteers were radically cured of malaria and during the next 126 days 25 of the 28 were reinfected. Resistance to malaria was not correlated with antibodies to malaria Ag, but was significantly correlated with lymphocyte responses to CS protein residues 361-380 and 371-390. Among the 25 volunteers who became re-infected with malaria, lymphocytes from only two responded to a peptide including residues 361-380 of the *P. falciparum* CS protein, and only one to peptide 371-390. In contrast, lymphocytes from all three volunteers who did not become infected responded to peptide 361-380 ($p = 0.003$), and lymphocytes from two of the three responded to peptide 371-390 ($p = 0.023$). The significant correlation between proliferation to peptides 361-380 and 371-390 and resistance to malaria suggests that at least one epitope within these overlapping peptides is involved in a protective cellular immune response. The data support inclusion of these residues in future CS protein vaccines.

Development of effective subunit vaccines requires identification of relevant epitopes and a method of immunization that induces an optimal immune response. The first human malaria vaccines have been designed to elicit antibodies to sporozoites, the stage of the parasite

transmitted to man by mosquitoes. A repetitive epitope on the CS¹ protein of *Plasmodium falciparum* was selected as the target antigen, because antibodies to the analogous epitope on a murine malaria CS protein protect mice against infection with sporozoites (1, 2) and antibodies to the *P. falciparum* repeat region mediate *in vitro* reactions thought to indicate protective immunity (3-6). However, the subunit sporozoite vaccines tested to date have been poorly immunogenic for humans, and only two of five volunteers who received the highest doses of these vaccines were protected against sporozoite-induced malaria (7, 8). One method for improving the antibody response to the CS repeat region would be to include additional Th epitopes in a vaccine. If these were derived from the CS protein, boosting of antibodies might occur after exposure to sporozoites (9, 10), and repeated doses of vaccine might not be required after primary immunization.

T cell epitopes are also important targets for the protective cell-mediated immunity that develops after immunization with irradiated sporozoites (2, 11-13). The effector arm of such immunity apparently requires lymphocytes of the suppressor/cytotoxic phenotype and IFN- γ (12, 13). The Ag responsible for this potent protective immune response are unknown. They may include T epitopes on the CS protein, because immunization of mice with an attenuated strain of *Salmonella typhimurium* transformed with the *P. berghei* CS gene induces protection against sporozoite challenge in the absence of anti-sporozoite antibodies (14).

Most adults in malaria endemic areas have antibodies to the repeat region of the CS protein (6, 15), but few have T lymphocytes sensitized to the same epitope (16-19), suggesting the existence of other Th epitopes (17, 19). These adults develop an immune response that renders them less susceptible to malaria than are children, and some adults less susceptible than others. If this immune response is directed against sporozoites, it is probably cellular in nature, because naturally acquired antibodies to sporozoites do not prevent human malaria infection (15). In the present study we identified non-repeat region, CS protein T epitopes that may provide help for produc-

Received for publication July 29, 1988.

Accepted for publication November 22, 1988.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹S.L.H. was supported in part by the Naval Medical Research and Development Command Work Unit No. 3M0102BS10AKIII. The opinions and assertions herein are those of the authors and are not to be construed as official or as reflecting the views of the Navy Department or the naval service at large.

*Address correspondence and reprint requests to Dr. Stephen L. Hoffman, Infectious Diseases Department, Naval Medical Research Institute Annex, 12300 Washington Ave., Rockville, MD 20852.

²Abbreviations used in this paper: CS, circumsporozoite protein; R32LR, a protein synthesized in *Escherichia coli*, that included 32 tetrapeptides derived from the *P. falciparum* CS protein; MDP, gNANP; NVDP, LR; peptides 361-380 and 371-390, synthetic peptide corresponding to residues 361-380 and 371-390 of the deduced amino acid sequence of the 768 clone of *P. falciparum*; SI, stimulation index.

TABLE I
Lymphocyte proliferative response to CS protein Ag and level of antibodies to the CS protein repeat region in 11 individuals whose lymphocytes proliferated after stimulation with at least one peptide

Antibody to CS Protein ^a	Stimulation Indices to each Peptide in Positive Responders						
	36-55	101-120	R32LR	326-345	351-370	361-380	371-390
0.09							5.58
0.34				30.21			
0.34				6.64		7.85	
0.39 ^b						5.85	6.49
0.44	4.32	2.32					
0.56					3.55		
0.77 ^b						14.65	12.42
0.82					2.99		4.39
1.07 ^b			5.4			15.57	
1.42						5.58	
1.77			6				

^a The absorbance (414 nm) of sera at a dilution of 1/100.

^b These three individuals did not develop recurrent malaria during the 126-day observation period.

Figure 2. Cumulative incidence of *P. falciparum* infections after radical cure and the entomologic inoculation rate during the 126 days of the study. The values for entomologic inoculation rate represent the mean entomologic inoculation rate during the preceding 14 days.




TABLE II
Comparison of day 0 lymphocyte proliferative responses to stimulation with peptides 361-380 and 371-390 in the 25 volunteers who did and the three who did not develop recurrent parasitemia^a

Developed Recurrent parasitemia	Stimulation with Peptides 361-380 and 371-390				
	N	No. Positive		SI	
		361-380	371-390	361-380	371-390
Yes	25	2	1	1.5 ± 1.90	1.3 ± 1.38
No	3	3	2	12.0 ± 5.37	6.7 ± 5.69

^a The results are expressed as the number of volunteers with a positive proliferative response, and the mean ± SD SI of the entire group. The group that did not develop recurrent parasitemia had a significantly greater proportion of responders to peptides 361-380 and 371-390 ($p = 0.003$ and 0.023 , respectively, Fisher's exact test, one-tailed), and a significantly greater mean SI ($p < 0.0005$, Student's t -test, two-tailed) than did the group that did develop recurrent parasitemia.

DISCUSSION

These studies delineate non-repeat region T epitopes on the CS protein of *P. falciparum*, suggest that some of these sites are involved in providing help for production of antibodies to the repeat region, and for the first time identify a region of the CS protein that may be involved in a protective cellular immune response.

Most individuals failed to respond to each peptide. This was apparently not due to malaria-related immuno-

suppression, or to an inappropriate in vitro dose of Ag, because repeating the assays 21 days after radical cure of malaria and testing with Ag concentrations from 0.23 to 60 $\mu\text{g/ml}$ did not significantly alter the results (C. F. Mason, manuscript in preparation). It may reflect genetic restriction of human response to T epitopes of the *P. falciparum* CS protein (19), a phenomenon demonstrated in mice for two regions of the *P. falciparum* CS protein (9, 23, 24), or the masking of the response to poor T epitopes on the CS protein by more dominant ones on the sporozoite, as has been shown for lysozyme (25). Despite the poor response to all the peptides, most individuals made antibodies to the repeat region. T cell help for production of antibodies to the repeats could have been provided by non-CS protein Th cell epitopes on sporozoites. However, the data suggest that a repeat region epitope and non-repeat region T cell epitopes on the CS protein, specifically those included within residues 351-380, provide T cell help for repeat antibodies. This interpretation is consistent with our finding that there are adequate Th cell sites on the *P. berghei* CS protein to completely overcome genetic restriction of response to the predominant *P. berghei* CS protein repeat.⁴

⁴ Hoffman, S. L., J. A. Berzofsky, D. Isenbarger, E. Zeltzer, W. R. Majarian, M. Gross, and W. R. Ballou. Immune response gene regulation of immunity to *Plasmodium berghei* sporozoites and circumsporozoite protein vaccines: Overcoming genetic restriction with whole organism and subunit vaccines. Submitted for publication.

Perhaps, the most important finding was the correlation between proliferation to peptides 361-380 (IKPGS-ANKPKDEL DYENDIEC) and 371-390 (DEL DYENDIEK-KICKMEK CSC) and resistance to re-infection with malaria (Table II). The three individuals who did not develop recurrent malaria and the 25 individuals who did had similar levels of antibodies to the repeat region of the CS protein, consistent with our previous findings, that in this village naturally acquired antibodies to sporozoites do not protect against malaria (15). Individuals in the two groups also had similar levels of antibodies to asexual, blood stages of *P. falciparum*. The significant difference between the non-infected and infected individuals in T cell proliferative responses to peptides 361-380 and 371-390 suggests that the proliferation reflects a protective cellular immune response. It is intriguing to speculate that one or more epitopes included within residues 361-390 are involved directly, or indirectly via Th cells, in a CD8⁺ cytotoxic T cell-mediated protective immune response,⁵ analogous to that demonstrated in the murine malaria models (12, 13). This can only be proven by immunization and challenge studies in human volunteers. However, this hypothesis is supported by preliminary experiments with an attenuated strain of *S. typhimurium* transformed with portions of the *P. berghei* CS gene. The carboxyl-terminal portion of the *P. berghei* CS protein, containing the region analogous to *P. falciparum* residues 361-390, appears to be required for induction of protective immunity (W. R. Ballou, personal communication).

Peptides 361-380 and 371-390 include residues known to vary among strains of *P. falciparum* (26). These peptides were derived from a Brazilian strain, and the three protected individuals were exposed to more than 100 infective bites in East Africa. If the response to these peptides reflects a protective immune response, either Kenyan and Brazilian parasites are conserved at the relevant site(s) along this sequence, or the protective response associated with these antigens is not restricted by minor amino acid substitutions.

The identification of a region of the *P. falciparum* CS protein that includes at least one T cell site that may be involved in a protective cellular immune response supports including this region in future malaria vaccines. However, only 21% of the individuals in this study responded to peptides 361-380 or 371-390, whereas all volunteers immunized with large numbers of irradiated sporozoites were protected against sporozoite challenge (27-30). This suggests that there are other protective T cell sites on the sporozoite or exoerythrocytic stages of *P. falciparum*, or that experimental immunization with large numbers of sporozoites over a short period of time induces a more potent immune response than does long term natural exposure to small numbers of sporozoites. Major challenges to further vaccine development will be to determine if there are additional protective T sites, and to develop methods of immunization that induce the required T cell immunity.

⁵ Since completion of this work residues 368-390 have been found to include the only cytotoxic T cell site on the *P. falciparum* CS protein in H10 BR (H-2^b) mice (Kumar S., L. H. Miller, I. A. Quakyi, D. B. Keister, R. A. Houghten, W. L. Maloy, B. Moss, J. A. Berzofsky, and M. F. Good, 1988. Cytotoxic T cells specific for the circumsporozoite protein of *Plasmodium falciparum*. *Nature* 334:258).

Acknowledgments. The authors thank the Saradidi Rural Health Project, D. O. Kaseje, C. R. Roberts, and C. L. Diggs for support and encouragement; L. T. Cannon, J. Koros, M. Adoyo-Adoyo, J. Kenner, R. Gore, J. Williams, and R. Hajkowski, for technical assistance; M. F. Good for synthetic peptides; and G. Wasserman and J. F. Young for purified R32LR.

REFERENCES

- Potocnjak, P., N. Yoshida, and R. S. Nussenzweig. 1980. Monovalent fragments (Fab) of monoclonal antibodies to a sporozoite surface antigen (pb44) protect mice against malaria infection. *J. Exp. Med.* 151:1504.
- Egan, J. E., J. L. Weber, W. R. Ballou, M. R. Hollingdale, W. R. Majarian, D. M. Gordon, W. L. Maloy, S. L. Hoffman, R. A. Wirtz, I. Schneider, G. R. Woollett, J. F. Young, and W. T. Hockmeyer. 1987. Efficacy of murine malaria sporozoite vaccines: Implications for human vaccine development. *Science* 236:453.
- Ballou, W. R., J. Rothbard, R. A. Wirtz, et al. 1985. Immunogenicity of synthetic peptides from circumsporozoite protein of *Plasmodium falciparum*. *Science* 228:1436.
- Young, J. F., W. T. Hockmeyer, M. Gross, W. R. Ballou, R. A. Wirtz, J. H. Trosper, R. L. Beaudoin, M. R. Hollingdale, L. H. Miller, C. L. Diggs, and M. Rosenberg. 1985. Expression of *Plasmodium falciparum* circumsporozoite proteins in *Escherichia coli* for potential use in human malaria vaccine. *Science* 228:958.
- Zavala, F., J. P. Tam, A. H. Cochrane, I. Quakyi, R. S. Nussenzweig, and V. Nussenzweig. 1985. Rationale for development of a synthetic vaccine against *Plasmodium falciparum* malaria. *Science* 228:1436.
- Hoffman, S. L., R. Wistar, W. R. Ballou, M. R. Hollingdale, R. A. Wirtz, I. Schneider, H. A. Marwoto, and W. T. Hockmeyer. 1986. Immunity to malaria and naturally acquired antibodies to the circumsporozoite protein of *Plasmodium falciparum*. *N. Engl. J. Med.* 315:601.
- Ballou, W. R., S. L. Hoffman, J. A. Sherwood, M. R. Hollingdale, F. A. Neva, W. L. Hockmeyer, D. M. Gordon, I. Schneider, R. A. Wirtz, J. F. Young, G. F. Wasserman, P. Reeve, C. L. Diggs, and J. D. Chulay. 1987. Safety and efficacy of a recombinant DNA *Plasmodium falciparum* sporozoite vaccine. *Lancet* 1:1277.
- Herrington, D. A., D. F. Clyde, G. Losonsky, M. Cortesia, J. R. Murphy, J. Davis, S. Bagar, A. M. Felix, E. P. Heimer, D. Gillespie, E. Nardin, R. S. Nussenzweig, V. Nussenzweig, M. R. Hollingdale, and M. Levine. 1987. Safety and immunogenicity in man of a synthetic peptide malaria vaccine against *Plasmodium falciparum* sporozoites. *Nature* 328:257.
- Good, M. F., W. L. Maloy, M. N. Lunde, H. Margalit, J. L. Cornette, G. L. Smith, B. Moss, L. H. Miller, and J. A. Berzofsky. 1987. Construction of synthetic immunogen: use of a new T-helper epitope on malaria circumsporozoite protein. *Science* 235:1059.
- Hoffman, S. L., L. T. Cannon, J. A. Berzofsky, W. R. Majarian, J. F. Young, W. L. Maloy, and W. T. Hockmeyer. 1987. *Plasmodium falciparum*: sporozoite boosting of immunity due to a T-cell epitope on a sporozoite vaccine. *Exp. Parasitol.* 64:64.
- Chen, D. H., R. E. Tigelaar, and F. I. Weinbaum. 1977. Immunity to sporozoite-induced malaria infection in mice. I. The effect of immunization on T and B cell deficient mice. *J. Immunol.* 118:1322.
- Schofield, L. J., Willaquiran, A., Ferreira, H., Schellekens, R., Nussenzweig, and V. Nussenzweig. 1987. Gamma Interferon, CD8⁺ cells (cytotoxic/suppressors) are required for protection in mice immunized with malaria sporozoites. *Nature* 330:664.
- Weiss, W. R., M. Sedegah, R. L. Beaudoin, L. H. Miller, and M. F. Good. 1988. CD8⁺ cells (cytotoxic/suppressors) are required for protection in mice immunized with malaria sporozoites. *Proc. Natl. Acad. Sci. USA* 85:573.
- Sadoff, J. C., W. R. Ballou, L. S. Baron, W. R. Majarian, R. N. Brey, W. T. Hockmeyer, J. F. Young, S. J. Cryz, J. Ou, G. H. Lowell, and J. D. Chulay. 1988. Oral *Salmonella typhimurium* vaccine expressing circumsporozoite protein protects against malaria. *Science* 240:639.
- Hoffman, S. L., C. N. Oster, C. V. Plowe, G. R. Woollett, J. C. Beier, J. D. Chulay, R. A. Wirtz, M. R. Hollingdale, and M. Mugambi. 1987. Naturally acquired antibodies to sporozoites do not prevent malaria vaccine development. Implications. *Science* 237:639.
- Campbell, J. R., F. P. Paleologo, E. D. Franke, S. Ratiwanto, H. Hadiputranto, I. Kurniawan, R. Wistar Jr., S. L. Hoffman, B. J. Anis, and L. W. Laughlin. 1988. Immune response to the circumsporozoite protein of *Plasmodium falciparum*: T helper activity in humans is based on recognition of a T cell epitope located outside of the central repeat region. *Am. J. Trop. Med. Hyg.* 39:232.
- Good, M. F., D. Pombo, I. A. Quakyi, E. M. Riley, R. A. Houghten, A. Menon, D. W. Alling, J. A. Berzofsky, and L. H. Miller. 1988. Human T cell recognition of the circumsporozoite protein of *Plasmodium falciparum*. Immunodominant T cell domains map to the

- polymorphic regions of the molecule. *Proc. Natl. Acad. Sci. USA* 85:1199.
18. Webster, H. K., M. Ho, S. Loareesuwan, K. Pavanand, Y. Wattanagoon, D. A. Warrell, and W. T. Hockmeyer. 1988. Lymphocyte responsiveness to a candidate malaria sporozoite vaccine (R32tet32) of individuals with naturally acquired *Plasmodium falciparum* malaria. *Am. J. Trop. Med. Hyg.* 38:37.
 19. Sinigaglia F., M. Guttinger, D. Gillesse, D. M. Doran, B. Takacs, H. Matile, A. Trzeciak, and J. R. Pink. 1988. Epitopes recognized by human T lymphocytes on malaria circumsporozoite protein. *Eur. J. Immunol.* 18:633.
 20. Dame, J. B., J. L. Williams, T. F. McCutchan, J. L. Weber, R. A. Wirtz, W. T. Hockmeyer, W. L. Maloy, J. D. Haynes, I. Schneider, D. Roberts, G. S. Sanders, E. P. Reddy, C. L. Diggs, and L. H. Miller. 1984. Structure of the gene encoding the immunodominant surface antigen on the sporozoite of the human malaria parasite, *Plasmodium falciparum*. *Science* 225:593.
 21. Houghten, R. A. 1985. General method for the rapid solid-phase synthesis of large numbers of peptides: specificity of antigen-antibody interaction at the level of individual amino acids. *Proc. Natl. Acad. Sci. USA* 82:5131.
 22. DeLisi, C., and J. A. Berzofsky. 1985. T-cell antigenic sites tend to be amphipathic structures. *Proc. Natl. Acad. Sci. USA* 82:7048.
 23. Good, M. F., J. A. Berzofsky, W. L. Maloy, Y. Hagashi, N. Fujii, W. T. Hockmeyer, and L. H. Miller. 1986. Genetic control of the immune responses in mice to *Plasmodium falciparum* sporozoite vaccine. Widespread non-responsiveness to single malaria T-epitope in highly repetitive vaccine. *J. Exp. Med.* 164:655.
 24. Del Giudice, G., J. A. Cooper, J. Merino, A. S. Verdini, A. Pessi, A. R. Togna, H. D. Engers, G. Corradin, and P-H. Lambert. 1986. The antibody response in mice to carrier-free synthetic polymers of *Plasmodium falciparum* circumsporozoite repetitive epitope is I-A^b restricted: possible implications for malaria vaccines. *J. Immunol.* 137:2952.
 25. Gammon, G., N. Shastri, J. Cogswell, S. Wilbur, S. Sadegh-Nasseri, U. Krzych, A. Miller, and E. Sercarz. 1987. The choice of T-cell epitopes utilized on a protein antigen depends on multiple factors distant from, as well as at the determinant site. *Immunol. Rev.* 98:53.
 26. de la Cruz, V. F., A. A. Lal, and T. F. McCutchan. 1987. Sequence variation in putative functional domain of the circumsporozoite protein of *Plasmodium falciparum*. Implications for vaccine development. *J. Biol. Chem.* 262:11935.
 27. Clyde, D. F., V. C. McCarthy, R. M. Miller, and R. B. Hornick. 1973. Specificity of protection of man immunized against sporozoite-induced falciparum malaria. *Am. J. Med. Sci.* 266:398.
 28. Rieckmann, K. H., P. E. Carson, R. L. Beaudoin, J. S. Cassells, and K. W. Sell. 1974. Sporozoite-induced immunity in man against an Ethiopian strain of *Plasmodium falciparum*. *Trans. R. Soc. Trop. Med. Hyg.* 68:258.
 29. Clyde, D. F., V. C. McCarthy, R. M. Miller, and W. E. Woodward. 1975. Immunization of man against falciparum and vivax malaria by use of attenuated sporozoites. *Am. J. Trop. Med. Hyg.* 24:397.
 30. Rieckmann, K. H., R. L. Beaudoin, J. S. Cassells, and K. W. Sell. 1979. Use of attenuated sporozoites in the immunization of human volunteers against falciparum malaria. *Bull. WHO* 57 (Suppl. 1):261.



Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail. and/or Special
A-1	20